

Frailty predicts medication-related harm requiring healthcare: A UK multicentre prospective cohort study

Parekh N, Ali K, Stevenson JM, Davies JG, Schiff R, Harchowal J, Van der Cammen T, Rajkumar C

Introduction

Frailty has been under investigated as a risk factor for medication-related harm (MRH) in older adults[1]. We sought to determine whether frailty is independently associated with MRH in a large multicentre prospective cohort, the PRIME study.

Methods

The PRIME study recruited 1280 older adults at hospital discharge from 5 hospitals in England between 2013 to 2015[2]. MRH and associated healthcare use within 8-weeks post-discharge were identified by senior pharmacists using (1) hospital readmission data, (2) primary care records, (3) patient telephone interviews. Based on the Rockwood approach[3], we developed a frailty index including 55 deficits from multiple domains (morbidity, cognition, mood, strength and mobility, nutrition, daily function). Frailty was defined using the established cut-off of $\geq 20\%$ deficits[4], and internally validated using Kaplan-Meier plots comparing survival in frail and non-frail patients. We then used logistic regression analysis to investigate the relationship between frailty and MRH requiring healthcare.

Results

1116 patients completed follow-up (median age 81.9 years, range 65-103 years, 58.4% female). 446 patients (40%) were frail in our cohort. 36% of frail patients experienced MRH compared with 25% in non-frail patients. There was a strong relationship between frailty and MRH (OR 1.67, 95% CI 1.29-2.17, $p < 0.001$). A significant relationship between frailty and MRH remained on multivariable regression, adjusting for polypharmacy, age and gender (OR 1.37, 95% CI 1.04-1.81, $p = 0.027$). Frail patients had significantly reduced 18-month survival (Log-Rank test $p < 0.001$).

Key Conclusions

Frailty is a predictor of MRH requiring healthcare, independent of polypharmacy.

References

- [1] Parekh N, Ali K, Page A, Roper T, Rajkumar C. Incidence of medication-related harm in older adults following hospital discharge: a systematic review. *J Am Geriatr Soc* 2018;in press.
- [2] Stevenson J, Parekh N, Ali K, Timeyin J, Bremner S, Van Der Cammen T, et al. Protocol for a Prospective (P) study to develop a model to stratify the risk (RI) of medication (M) related harm in hospitalized elderly (E) patients in the UK (The PRIME study). *BMC Geriatr* 2016;16:22. doi:10.1186/s12877-016-0191-8.
- [3] Searle SD, Mitnitski A, Gahbauer EA, Gill TM, Rockwood K. A standard procedure for creating a frailty index. *BMC Geriatr* 2008;8:24. doi:10.1186/1471-2318-8-24.
- [4] Mitnitski A, Song X, Skoog I, Broe G, Cox JL, Grunfeld E, et al. Relative Fitness and Frailty of Elderly Men and Women in Developed Countries and Their Relationship

with Mortality. J Am Geriatr Soc 2005;53:2184–9.